

**OSM rabbit pAb****Cat#: orb771538 (Manual)**


For research use only. Not intended for diagnostic use.

|                                 |   |
|---------------------------------|---|
| <b>Product Name</b>             | OSM rabbit pAb  |
| <b>Host species</b>             | Rabbit  |
| <b>Applications</b>             | IHC;IF;ELISA  |
| <b>Species Cross-Reactivity</b> | Human;Rat;Mouse;  |
| <b>Recommended dilutions</b>    | IHC-p 1:50-200, ELISA 1:10000-20000   |
| <b>Immunogen</b>                | Synthetic peptide from human protein at AA range: 50-100  |
| <b>Specificity</b>              | The antibody detects endogenous OSM   |
| <b>Formulation</b>              | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..  |
| <b>Storage</b>                  | Store at -20°C. Avoid repeated freeze-thaw cycles.  |
| <b>Protein Name</b>             | Oncostatin-M (OSM)  |
| <b>Gene Name</b>                | OSM   |
| <b>Cellular localization</b>    | Secreted.   |
| <b>Purification</b>             | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Clonality</b>                | Polyclonal  |


|                                |                    |
|--------------------------------|--------------------|
| <b>Concentration</b>           | 1 mg/ml            |
| <b>Observed band</b>           |                    |
| <b>Human Gene ID</b>           | 5008               |
| <b>Human Swiss-Prot Number</b> | P13725             |
| <b>Alternative Names</b>       | Oncostatin-M (OSM) |

### Background


This gene encodes a member of the leukemia inhibitory factor/oncostatin-M (LIF/OSM) family of proteins. The encoded preproprotein is proteolytically processed to generate the mature protein. This protein is a secreted cytokine and growth regulator that inhibits the proliferation of a number of tumor cell lines. This protein also regulates the production of other cytokines, including interleukin 6, granulocyte-colony stimulating factor and granulocyte-macrophage colony stimulating factor in endothelial cells. This gene and the related gene, leukemia inhibitory factor, also present on chromosome 22, may have resulted from the duplication of a common ancestral gene. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016],




**Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200**



**Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200**



**Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200**



**Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200**