

**KAT6A rabbit pAb****Cat#: orb772089 (Manual)**

For research use only. Not intended for diagnostic use.

<b>Product Name</b>	KAT6A rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse;Rat
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 160-240
<b>Specificity</b>	KAT6A Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide..
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Histone acetyltransferase KAT6A (EC 2.3.1.48) (MOZ, YBF2/SAS3, SAS2 and TIP60 protein 3) (MYST-3) (Monocytic leukemia zinc finger protein) (Runt-related transcription factor-binding protein 2) (Zinc f
<b>Gene Name</b>	KAT6A MOZ MYST3 RUNXBP2 ZNF220
<b>Cellular localization</b>	Nucleus. Nucleus, nucleolus. Nucleus, nucleoplasm. Nucleus, PML body. Recruited into PML body after DNA damage.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

<b>Clonality</b>	Polyclonal
------------------	------------

<b>Concentration</b>	1 mg/ml
----------------------	---------

<b>Observed band</b>	220kD
----------------------	-------

<b>Human Gene ID</b>	7994
----------------------	------

<b>Human Swiss-Prot Number</b>	Q92794
--------------------------------	--------

<b>Alternative Names</b>	
--------------------------	--

### Background

This gene encodes a member of the MOZ, YBFR2, SAS2, TIP60 family of histone acetyltransferases. The protein is composed of a nuclear localization domain, a double C2H2 zinc finger domain that binds to acetylated histone tails, a histone acetyl-transferase domain, a glutamate/aspartate-rich region, and a serine- and methionine-rich transactivation domain. It is part of a complex that acetylates lysine-9 residues in histone 3, and in addition, it acts as a co-activator for several transcription factors. Allelic variants of this gene are associated with autosomal dominant mental retardation-32. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015],