

## www.biorbyt.com

## GCNT4 rabbit pAb


## Cat#: orb772319 (Manual)

For research use only. Not intended for diagnostic use.

Product Name	GCNT4 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at AA range: 60-140
Specificity	GCNT4 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N- acetylglucosaminyltransferase 4 (EC 2.4.1.102) (Core 2-branching enzyme 3) (Core2-GlcNAc-transferase 3) (C2GnT3)
Gene Name	GCNT4
Cellular localization	Golgi apparatus membrane ; Single-pass type II membrane protein .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Clonality	Polyclonal



Concentration	1 mg/ml
Observed band	49kD
Human Gene ID	51301
Human Swiss-Prot Number	Q9P109
Alternative Names	
Background	catalytic activity:UDP-N-acetyl-D-glucosamine + beta-D-galactosyl-1,3-N- acetyl-D-galactosaminyl-R = UDP + beta-D-galactosyl-1,3-(N-acetyl-beta- D-glucosaminyl-1,6)-N-acetyl-D-galactosaminyl- R.,function:Glycosyltransferase that mediates core 2 O-glycan branching, an important step in mucin-type biosynthesis. Does not have core 4 O-glycan or I-branching enzyme activity.,online information:Core 2 beta-1,6-N- acetylglucosaminyltransferase 3.,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 14 family.,tissue specificity:Predominantly expressed in thymus. Weakly expressed in pancreas, peripheral blood leukocytes, placenta, small intestine and stomach. Barely detectable in liver, spleen, lung and lymph node.,



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000