

**UNC5D rabbit pAb****Cat#: orb772768 (Manual)**

For research use only. Not intended for diagnostic use.

<b>Product Name</b>	UNC5D rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 160-240
<b>Specificity</b>	UNC5D Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide..
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Netrin receptor UNC5D (Protein unc-5 homolog 4) (Protein unc-5 homolog D)
<b>Gene Name</b>	UNC5D KIAA1777 UNC5H4 UNQ6012/PRO34692
<b>Cellular localization</b>	Cell membrane ; Single-pass type I membrane protein .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal

<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	104kD
<b>Human Gene ID</b>	137970
<b>Human Swiss-Prot Number</b>	Q6UXZ4
<b>Alternative Names</b>	

**Background**

function:Receptor for netrin. May be involved in axon guidance by mediating axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand.,PTM:Phosphorylated on cytoplasmic tyrosine residues.,PTM:Proteolytically cleaved by caspases during apoptosis. The cleavage does not take place when the receptor is associated with netrin ligand. Its cleavage by caspases is required to induce apoptosis.,similarity:Belongs to the unc-5 family.,similarity:Contains 1 death domain.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 ZU5 domain.,similarity:Contains 2 TSP type-1 domains.,subunit:Interacts with the cytoplasmic part of DCC.,