

KDM3B rabbit pAb**Cat#: orb773468 (Manual)**

For research use only. Not intended for diagnostic use.

Product Name	KDM3B rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at AA range: 230-310
Specificity	KDM3B Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Lysine-specific demethylase 3B (EC 1.14.11.-) (JmjC domain-containing histone demethylation protein 2B) (Jumonji domain-containing protein 1B) (Nuclear protein 5qNCA)
Gene Name	KDM3B C5orf7 JHDM2B JMJD1B KIAA1082
Cellular localization	Nucleus .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal

Concentration	1 mg/ml
Observed band	193kD
Human Gene ID	51780
Human Swiss-Prot Number	Q7LBC6
Alternative Names	

Background

cofactor: Binds 1 Fe(2+) ion per subunit., domain: Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs are known to mediate the association with nuclear receptors., function: Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Demethylation of Lys residue generates formaldehyde and succinate. May have tumor suppressor activity., miscellaneous: Its gene is located in the 5q region of the genome which is deleted in del(5q) interstitial deletion, a frequent deletion found in myeloid leukemias and myelodysplasias, suggesting that it may be a good candidate for the del(5q) tumor suppressor gene., similarity: Belongs to the JHDM2 histone demethylase family., similarity: Contains 1 JmjC domain., tissue specificity: Ubiquitous. Highly expressed in placenta, skeletal muscle, kidney, heart and liver.,