



HIPK3 rabbit pAb

Cat#: orb773525 (Manual)

For research use only. Not intended for diagnostic use.

Product Name HIPK3 rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein . at AA range: 770-850

HIPK3 Polyclonal Antibody detects endogenous levels of protein. **Specificity**

Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Homeodomain-interacting protein kinase 3 (EC 2.7.11.1) (Androgen receptor-interacting nuclear protein kinase) (ANPK) (Fas-interacting **Protein Name**

serine/threonine-protein kinase) (FIST) (Homolog of protein kinase)

HIPK3 DYRK6 FIST3 PKY

Cytoplasm . Nucleus . Cellular localization

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal

Gene Name





Concentration 1 mg/ml

Observed band 133kD

Human Gene ID 10114

Human Swiss-Prot Number Q9H422

Alternative Names

Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Seems to negatively regulate apoptosis by promoting FADD phosphorylation. Enhances androgen receptor-mediated transcription. May act as a transcriptional corepressor for NK homeodomain transcription factors.,PTM:Autophosphorylated, but autophosphorylation is not required for catalytic activity.,PTM:May be sumoylated.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. HIPK subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with Nkx1-2. Interacts with FAS and DAXX. Probably part of a complex consisting of HIPK3, FAS and FADD. Interacts with and stabilizes ligand-bound androgen receptor (AR) (By similarity). Interacts with UBL1/SUMO-1.,tissue specificity:Overexpressed in multidrug resistant cells. Highly expressed in heart and skeletal muscle, and at lower levels in placenta, pancreas, brain, spleen, prostate, thymus, testis, small intestine, colon and leukocytes. Not found in liver and lung.,