



## SGK3 rabbit pAb

Cat#: orb773704 (Manual)

For research use only. Not intended for diagnostic use.

**Product Name** SGK3 rabbit pAb

**Host species** Rabbit

**Applications** WB;ELISA

**Species Cross-Reactivity** Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from part region of human protein

SGK3 Polyclonal Antibody detects endogenous levels of protein. **Specificity** 

**Formulation** Liquid in PBS containing 50% glycerol, and 0.02% sodium azide..

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Serine/threonine-protein kinase Sgk3 (EC 2.7.11.1) (Cytokine-independent survival kinase) (Serum/glucocorticoid-regulated kinase 3) **Protein Name** 

(Serum/glucocorticoid-regulated kinase-like)

Gene Name SGK3 CISK SGKL

Cellular localization Cytoplasmic vesicle . Early endosome. Recycling endosome. Endosomal

localization is a prerequisite for complete kinase activity. It is essential for its colocalization with the kinase responsible for phosphorylating Ser-486 thus allowing PDPK1 phosphorylation of Thr-320 resulting in complete activation of SGK3. Localized in vesicle-like structures and in the early endosome. Colocalizes with SLC9A3/NHE3 in the recycling endosomes.



## www.biorbyt.com

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Purification

**Clonality** Polyclonal

Concentration 1 mg/ml

**Observed band** 54kD

**Human Gene ID** 100533105

**Human Swiss-Prot Number** Q96BR1

**Alternative Names** 

**Background** 

This gene is a member of the Ser/Thr protein kinase family and encodes a phosphoprotein with a PX (phox homology) domain. The protein phosphorylates several target proteins and has a role in neutral amino acid transport and activation of potassium and chloride channels. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],