

## **L-galactono-1, 4-lactone Dehydrogenase**

### **Microplate Assay Kit**

**Cat #: orb390756 (manual)**

Detection and Quantification of L-galactono-1, 4-lactone Dehydrogenase (GalLDH) Activity in Tissue extracts, Cell lysate Samples.

*For research use only. Not for diagnostic or therapeutic procedures.*

## INTRODUCTION

L-galactono-1, 4-lactone Dehydrogenase (EC 1.3.2.3) catalyzes the last step in the main pathway of vitamin C (L-ascorbic acid) biosynthesis in higher plants.

The enzyme catalysated reaction products reduced Cyt c can be measured at a colorimetric readout at 550 nm.

## KIT COMPONENTS

Component	Volume	Storage
96-Well Microplate	1 plate	
Assay Buffer	30 ml x 4	4 °C
Substrate I	Powder x 1	4 °C, keep in dark
Substrate II	Powder x 1	4 °C
Technical Manual	1 Manual	

### Note:

**Substrate I:** add 17 ml distilled water to dissolve before use.

**Substrate II:** add 1 ml distilled water to dissolve before use.

## MATERIALS REQUIRED BUT NOT PROVIDED

1. Microplate reader to read absorbance at 550 nm
2. Distilled water
3. Pipettor, multi-channel pipettor
4. Pipette tips
5. Mortar
6. Centrifuge
7. Timer
8. Ice

## SAMPLE PREPARATION

### 1. For cell and bacteria samples

Collect cell or bacteria into centrifuge tube, discard the supernatant after centrifugation, add 1 ml Assay buffer for  $5 \times 10^6$  cell or bacteria, sonicate (with power 20%, sonicate 3s, interval 10s, repeat 30 times); centrifuged at 13000g 4 °C for 10 minutes, take the supernatant into a new centrifuge tube and keep it on ice for detection.

### 2. For tissue samples

Weigh out 0.1 g tissue, homogenize with 1 ml Assay buffer on ice, centrifuged at 13000g 4 °C for 10 minutes, take the supernatant into a new centrifuge tube and keep it on ice for detection.

## ASSAY PROCEDURE

Warm the Substrate I and Substrate II to room temperature before use.

Add following reagents in the microplate:

Reagent	Sample	Blank
Sample	20 $\mu$ l	--
Distilled water	--	20 $\mu$ l
Substrate I	170 $\mu$ l	170 $\mu$ l
Substrate II	10 $\mu$ l	10 $\mu$ l
Mix, measured at 550 nm and record the absorbance of 10th second and 130th second.		

## CALCULATION

**Unit Definition:** One unit of GalLDH is the amount of enzyme that will reduce 1  $\mu$ mol Cyt c per minute.

1. According to the protein concentration of sample

$$\begin{aligned} \text{GalLDH (U/mg)} &= [ (\text{OD}_{\text{Sample (130S)}} - \text{OD}_{\text{Sample (10S)}}) - (\text{OD}_{\text{Blank (130S)}} - \text{OD}_{\text{Blank (10S)}}) ] / (\epsilon \times d) \times V_{\text{Total}} \times 10^6 / \\ &\quad (V_{\text{Sample}} \times C_{\text{Protein}}) / T \\ &= 481.7 \times [ (\text{OD}_{\text{Sample (130S)}} - \text{OD}_{\text{Sample (10S)}}) - (\text{OD}_{\text{Blank (130S)}} - \text{OD}_{\text{Blank (10S)}}) ] / C_{\text{Protein}} \end{aligned}$$

2. According to the weight of sample

$$\begin{aligned} \text{GalLDH (U/g)} &= [ (\text{OD}_{\text{Sample (130S)}} - \text{OD}_{\text{Sample (10S)}}) - (\text{OD}_{\text{Blank (130S)}} - \text{OD}_{\text{Blank (10S)}}) ] / (\epsilon \times d) \times V_{\text{Total}} \times 10^6 / (W \\ &\quad \times V_{\text{Sample}} / V_{\text{Assay}}) / T \\ &= 481.7 \times [ (\text{OD}_{\text{Sample (130S)}} - \text{OD}_{\text{Sample (10S)}}) - (\text{OD}_{\text{Blank (130S)}} - \text{OD}_{\text{Blank (10S)}}) ] / W \end{aligned}$$

3. According to the quantity of cells or bacteria

$$\begin{aligned} \text{GalLDH (U/10}^4) &= [ (\text{OD}_{\text{Sample (130S)}} - \text{OD}_{\text{Sample (10S)}}) - (\text{OD}_{\text{Blank (130S)}} - \text{OD}_{\text{Blank (10S)}}) ] / (\epsilon \times d) \times V_{\text{Total}} \times 10^6 / (N \\ &\quad \times V_{\text{Sample}} / V_{\text{Assay}}) / T \\ &= 481.7 \times [ (\text{OD}_{\text{Sample (130S)}} - \text{OD}_{\text{Sample (10S)}}) / (\text{OD}_{\text{Standard (130S)}} - \text{OD}_{\text{Standard (10S)}}) ] / N \end{aligned}$$

$\epsilon$ : molar extinction coefficient,  $17.3 \times 10^3$  L/mol/cm;

d: the optical path of 96-Well microplate, 0.6 cm;

$C_{\text{Protein}}$ : the protein concentration, mg/ml;

W: the weight of sample, g;

N: the quantity of cell or bacteria,  $N \times 10^4$ ;

$V_{\text{Total}}$ : the total volume of the enzymatic reaction, 0.2 ml;

$V_{\text{Sample}}$ : the volume of sample, 0.02 ml;

$V_{\text{Assay}}$ : the volume of Assay buffer, 1 ml;

T: the reaction time, 2 minutes.