

## **Product Datasheet**

## Rat Cluster of Differentiation 8 (CD8) ELISA Kit (orb1146831)

## **Description**

The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Cluster ofDifferentiation 8(CD8). Standards or samples are added to the appropriate microtiter plate wells then with a biotin-conjugated antibody specific to Cluster ofDifferentiation 8(CD8). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Cluster ofDifferentiation 8(CD8), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm  $\pm$  10nm. The concentration of Cluster ofDifferentiation 8(CD8) in the samples is then determined by comparing the OD of the samples to the standard curve.

**Reactivity** Rat

**Range** 0.16-10 ng/mL

**Concentration** 10 ng/mL

**Note** For research use only

## **Application notes**

standard: 10 ng/mL. Test principle: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Rat CD8. Standards or samples are added to the appropriate microtiter plate wells then with a biotin-conjugated antibody specific to Rat CD8. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Rat CD8, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of  $450 \, \mathrm{nm} \pm 10 \, \mathrm{nm}$ . The concentration of Rat CD8 in the samples is then determined by comparing the OD of the samples to the standard curve





Sample Types tissue homogenates, cell lysates, cell culture supernates and other biological


fluids

**Assay Time** 3.5h

Uniprot ID P07725

**Sensitivity** 0.062 ng/mL

**Expiration Date** Please enquire.



 $\begin{aligned} & \text{Email: } \underline{info@biorbyt.com}, \ \underline{support@biorbyt.com} \\ & \text{Phone: } \underline{+1~(415)~906-5211} \mid \text{Fax: } \underline{+1~(415)~651-8558} \end{aligned}$