

Product Datasheet

F(ab')₂ Fragment Goat Anti-Mouse IgG H&L, HRP conjugated (orb1151905)

Catalog Number	orb1151905
Category	Antibodies
Description	The F(ab') ₂ Fragment Goat Anti-Mouse IgG H&L, HRP conjugated is an HRP conjugated polyclonal antibody suitable for ELISA, IHC-Fr, IHC-P, WB. This product reacts with Mouse samples.
Clonality	Polyclonal
Species/Host	Goat
Isotype	IgG
Conjugation	HRP
Reactivity	Mouse
Form/Appearance	Liquid
Concentration	1.0 mg/ml
Buffer/Preservatives	10 mM TBS (pH=7.4) with 1% rAlbumin, 0.03% Proclin300 and 50% glycerol.
Purification	Affinity purified by Protein G
Immunogen	Native Mouse IgG
Tested applications	ELISA, IHC-Fr, IHC-P, WB
Dilution range	IHC-P=1:1000, WB=1:1000-20000, IHC-F=1:100-1000, ELISA=1:1000-20000
Antibody Type	Secondary Antibody

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom


Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Storage	Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Note	For research use only
Expiration Date	12 months from date of receipt.



Paraformaldehyde-fixed, paraffin embedded (rat cerebellum), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30 min, Antibody incubation with (S100B) Monoclonal Antibody, Unconjugated (orb500644) at 1:200 overnight at 4°C, An HRP-conjugated secondary (orb1151905, 1:300 dilution) was used for 20min at 37°C, DAB was used as the chromogen.



Paraformaldehyde-fixed, paraffin embedded (rat pancreas), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30 min, Antibody incubation with (GLP-1 (1G9)) Monoclonal Antibody, Unconjugated (orb10721) at 1:200 overnight at 4°C, An HRP-conjugated secondary (orb1151905, 1:300 dilution) was used for 20min at 37°C, DAB was used as the chromogen.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558