

Product Datasheet

NPHS1 Antibody (orb1239664)

Catalog Number	orb1239664
Category	Antibodies
Description	NPHS1 Antibody
Target	NPHS1
Clonality	Polyclonal
Species/Host	Rabbit
Isotype	IgG
Conjugation	Unconjugated
Reactivity	Human, Mouse, Rat
Form/Appearance	Liquid
Concentration	1 mg/mL
Buffer/Preservatives	Nephrin Antibody is supplied in PBS containing 0.02% sodium azide.
Purification	Nephrin Antibody is affinity chromatography purified via peptide column.
Immunogen	Anti-Nephrin antibody (orb1239664) was raised against a peptide corresponding to 14 amino acids near the carboxy terminus of human Nephrin. The immunogen is located within the last 50 amino acids of Nephrin.
UniProt ID	O60500
Tested applications	ELISA, IF, IHC-P, WB
Antibody Type	Primary Antibody

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States


Email: info@biorbyt.com, support@biorbyt.com

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Modifications	None
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Note	For research use only
NCBI	NP_004637
Expiration Date	12 months from date of receipt.



Western Blot Validation in Mouse Kidney Tissue Lysate with the (A) absence or the (B) presence of blocking peptide. Loading: 15 μ g of lysates per lane. Antibodies: Nephrin orb1239664 (1 μ g/mL), 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.



Western Blot Validation in Human and Mouse Cell Lines. Loading: 15 μ g of lysates per lane. Antibodies: Nephrin orb1239664 (2 μ g/mL), 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com


Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com


Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Immunohistochemistry Validation of Nephrin in Mouse Kidney Tissue. Immunohistochemical analysis of paraffin-embedded mouse kidney tissue using anti- Nephrin antibody (orb1239664) at 1 $\mu\text{g}/\text{ml}$. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4°C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.



Immunofluorescence Validation of Nephrin in Mouse Kidney Tissue. Immunofluorescent analysis of 4% paraformaldehyde-fixed mouse kidney cells labeling Nephrin with orb1239664 at 10 $\mu\text{g}/\text{mL}$, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (red).



Immunohistochemistry Validation of Nephrin in Rat Kidney Tissue. Immunohistochemical analysis of paraffin-embedded rat kidney tissue using anti- Nephrin antibody (orb1239664) at 5 $\mu\text{g}/\text{ml}$. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4°C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com


Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com


Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)



Immunofluorescence Validation of Nephrin in Rat Kidney Tissue. Immunofluorescent analysis of 4% paraformaldehyde-fixed rat kidney tissue labeling Nephrin with orb1239664 at 20 µg/mL, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (red) and DAPI staining (blue).



Apoptosis Assay Validation of Nephrin in Mouse Glomerulus (Chen et al., 2017). Glomerular cells of *Atgl* (-/-) mice were double labeled with TUNEL staining (dark brown nucleus indicated by red arrows) and immunofluorescence staining of nephrin detected by anti-nephin antibodies (orb1239664) (pink cytoplasm indicated by green arrows) as a marker for podocytes. Colocalization of TUNEL-positive cells and nephrin proved that apoptosis was induced in *Atgl* (-/-) mice as compared to WT mice.



Immunolocalization Validation of Nephrin in Human Placenta (Yun et al., 2015). Immunofluorescence staining showed Nephrin expression detected by anti-nephin antibodies (orb1239664) was clearly localized in villi (A-C) and fetal membranes, Amnion (J-L) and Chorion (M-O). The staining was markedly positive at apical membrane of villi (arrows in B and C) and amnion (arrow in L), and in the stromal cells of chorion (small arrow in O).

Biorbyt Ltd.


7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States


Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Immunofluorescence Validation of Nephrin in Mouse Podocyte (Li et al, 2013). Double immunofluorescence analysis of podocytic membrane protein nephrin (red) and nuclei stained with DAPI (blue). The presence of high glucose (HG) and neutralizing antibody (NtAb) which blocked epithelial growth factor (EGF) decreased nephrin expression while mesenchymal stem cells-conditioned medium (MSCs-CM) and recombinant human EGF (rhEGF) prevented the effect.



Induced Expression of Nephrin by Curcumin Treatment in the Renal Tissues of Type 1 Diabetic Rats (Soetikno et al., 2013). Nephrin expression detected by anti-nephin antibodies in type 1 diabetic rats. Nephrin was down-regulated in the vehicle-treated diabetic rats as compared to the control nondiabetic rats. However, this decrease in nephrin protein expression was markedly increased by curcumin treatment (P .05) to near-normal levels. (n = 5 in each group).



Immunofluorescence and Localization Validation of Nephrin in Cultured Rat Podocytes (Piwkowska et al., 2012). Immunofluorescence staining showed Nephrin expression (green) detected by anti-nephin antibodies and PKG1alpha (red). The co-localization of two antibodies (yellow) in rat podocytes was observed particularly at the tips of the cell processes.

Biorbyt Ltd.


7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States



Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



WB Validation of Nephrin in Glomeruli of Zucker Obese (ZO) and Zucker Lean (ZL) Rats (Piwkowska et al., 2013). The expression of nephrin detected by anti-nephrin antibodies did not change in ZO rats as compared to the control rats.



Immunohistochemistry Validation of Nephrin in Mouse Kidney (Toyama et al., 2012). Protein analysis for nephrin by immunohistochemistry with anti-nephrin antibodies in the kidney of wild-type or AMPD2-deficient mice at 2, 12 or 24 weeks of age. No difference between wild-type and AMPD2-deficient mice at any age was observed.



Regulated Expression Validation of Nephrin in Mouse Podocyte Cells Cultured in Normal Glucose (NG) Medium or High Glucose (HG) Medium (Huang et al., 2019). Western Blot analysis was used to assess the protein expression level of nephrin with anti-nephrin antibodies. Nephrin expression was down-regulated by PEGF treatment (NGP or HGP), which was reversed by the addition of C3 transferase.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558