

Product Datasheet

CCL27 Antibody (orb1272644)

Catalog Number	orb1272644
Category	Antibodies
Description	CCL27 Antibody
Target	CCL27
Clonality	Polyclonal
Species/Host	Rabbit
Conjugation	Unconjugated
Reactivity	Human
Form/Appearance	Lyophilized
Concentration	batch dependent
Immunogen	Produced from sera of rabbits immunized with highly pure recombinant Human CTACK. Human CTACK specific antibody was purified by affinity chromatography employing an immobilized Human CTACK matrix.
UniProt ID	Q5VZ77
Tested applications	ELISA, WB
Antibody Type	Primary Antibody
Modifications	None
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)


Note For research use only

NCBI [Q5VZ77](#)


Expiration Date 12 months from date of receipt.



To detect Human CTACK by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human CTACK (orb1272633) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human CTACK.



To detect Human CTACK by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 ug/ml. When used in conjunction with compatible secondary reagents the detection limit for recombinant Human CTACK is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.



To detect Human CTACK by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 ug/ml. When used in conjunction with compatible secondary reagents the detection limit for recombinant Human CTACK is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Biorbyt Ltd.


7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



This antibody stained HeLa cells. The primary antibody was incubated at 3.0 ug/ml overnight at 4 °C followed by a fluorescent labeled secondary antibody. Optimal concentrations and conditions may vary.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)