



Product Datasheet PRMT6 Antibody (orb1474396)

Catalog Number orb1474396

Category Antibodies

Description The PRMT6 Antibody is suitable for IF, IHC, WB. It is a Monoclonal, Unconjugated

antibody which raised against Purified recombinant human PRMT6 protein fragments expressed in E.coli. Purification: The antibody was purified by

immunogen affinity chromatography.

Clonality Monoclonal

Species/Host Mouse

Conjugation Unconjugated

Reactivity Human

Form/Appearance Liquid in PBS containing 50% glycerol, 0.5% rAlbumin and 0.02% sodium azide,

pH 7.3.

Purification The antibody was purified by immunogen affinity chromatography.

Immunogen Purified recombinant human PRMT6 protein fragments expressed in E.coli.

UniProt ID Q96LA8

Tested applications IF, IHC, WB

Dilution range WB (1/500 - 1/1000), IH (1/50 - 1/100), IF/IC (1/50 - 1/100)

Antibody Type Primary Antibody

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only







Entrez 55170

Expiration Date


12 months from date of receipt.



Western blot analysis of PRMT6 expression in Hela (A), 293T (B) whole cell lysates. (Predicted band size: 42 kD; Observed band size: 42 kD)



Immunohistochemical analysis of PRMT6 staining in human breast carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.29). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of PRMT6 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.