

Product Datasheet

PARK7/DJ1 Antibody (monoclonal, 4B10) (orb1474883)

Catalog Number orb1474883

Category Antibodies

Description Anti-PARK7/DJ1 Antibody (monoclonal, 4B10). Tested in Flow Cytometry, IF, IHC,

ICC, WB applications. This antibody reacts with Human.

Clonality Monoclonal

Species/Host Mouse

Isotype IgG2b

Conjugation Unconjugated

Reactivity Human

Form/Appearance Lyophilized

Concentration Adding 0.2 ml of distilled water will yield a concentration of 500 μg/ml.

Purification Immunogen affinity purified.

Immunogen E.coli-derived human PARK7 recombinant protein (Position: A2-D189). Human

PARK7 shares 91% amino acid (aa) sequence identity with both mouse and rat

PARK7.

UniProt ID Q99497

MW 22 kDa

Tested applications FC, ICC, IF, IHC, WB

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713

NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Biorbyt.com

Application notes Western blot, 0.25-0.5 μg/ml, Human Immunohistochemistry(Paraffin-embedded

Section), 2-5 μ g/ml, Human Immunocytochemistry/Immunofluorescence, 5 μ g/ml, Human Flow Cytometry (Fixed), 1-3 μ g/1x106 cells, Human. Adding 0.2 ml

of distilled water will yield a concentration of 500 μg/ml

Cross Reactivity No cross-reactivity with other proteins.


Antibody Type Primary Antibody

Clone Number 4B10

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only



Flow Cytometry analysis of Hela cells using anti-PARK7/DJ1 antibody. Overlay histogram showing Hela cells (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-PARK7/DJ1 Antibody (1 $\mu g/1x10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (5-10 $\mu g/1x10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu g/1x10^6$) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: <u>+44 (0)1223 859353</u> | Fax: <u>+1 (415) 651-8558</u>


Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States


Email: $\underline{info@biorbyt.com}$, $\underline{support@biorbyt.com}$ Phone: $\underline{+1 (415) 906-5211}$ | Fax: $\underline{+1 (415) 651-8558}$








IF analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody. PARK7/DJ1 was detected in an immunocytochemical section of Hela cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 μg/mL mouse anti-PARK7/DJ1 Antibody overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody. PARK7/DJ1 was detected in a paraffin-embedded section of human endometrial carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-PARK7/DJ1 Antibody overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody. PARK7/DJ1 was detected in a paraffin-embedded section of human esophageal squamous carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μg/ml mouse anti-PARK7/DJ1 Antibody overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States


Email: $\underline{info@biorbyt.com}$, $\underline{support@biorbyt.com}$ Phone: $\underline{+1 (415) 906-5211}$ | Fax: $\underline{+1 (415) 651-8558}$







IHC analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody. PARK7/DJ1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-PARK7/DJ1 Antibody overnight at 4°C. Peroxidase Conjugated Goat Antimouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



Western blot analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human U20S whole cell lysates, Lane 2: human HepG2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-PARK7/DJ1 antigen affinity purified monoclonal antibody at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PARK7/DJ1 at approximately 22 kDa. The expected band size for PARK7/DJ1 is at 22 kDa.

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: $\underline{info@biorbyt.com}$, $\underline{support@biorbyt.com}$ Phone: $\underline{+1 (415) 906-5211}$ | Fax: $\underline{+1 (415) 651-8558}$