

Product Datasheet

GRP78 Antibody: APC (orb148007)

Catalog Number orb148007

Description Mouse monoclonal to GRP78 (APC). GRP78 is a ubiquitously expressed, 78-kDa

glucose- regulated protein, and is commonly referred to as an immunoglobin chain binding protein (BiP). The BiP proteins are categorized as stress response proteins because they play an important role in the proper folding and assembly

of nascent protein and in the scavenging of misfolded proteins in the endoplasmic reticulum lumen. Translation of BiP is directed by an internal ribosomal entry site (IRES) in the 5' non-translated region of the BiP mRNA. BiP

IRES activity increases when cells are heat stressed. GRP78 is also critical for maintenance of cell homeostasis and the prevention of apoptosis. Lou et al. have provided findings that suggest GRP78 is essential for embryonic cell growth and pluripotent cell survival. In terms of diseases, GRP78 has been shown to be a reliable biomarker of hypoglycemia, to serve a neuroprotective function in neurons exposed to glutamate and oxidative stress, and its protein levels are

protein that results in severe glucose and oxygen deprivation could possible lead

reduced in the brains of Alzheimers patients. Also, the induction of the GRP78

Species/Host Mouse

Reactivity Bovine, Frog, Fungi, Hamster, Human, Monkey, Mouse, Plant, Rabbit, Rat

to drug resistance to anti-tumor drugs ..

Conjugation APC

Tested Applications ICC, IF, IHC

Immunogen His-tagged human GRP78

Target GRP78

Preservatives 95.64mM Phosphate, 2.48mM MES and 2mM EDTA

Concentration 1 mg/ml

Storage Conjugated antibodies should be stored according to the product label





Note For research use only

Application notes 0.5 μ g/ml of SMC-195 was sufficient for detection of Grp78 in 10 μ g of rat tissue

lysate by ECL immunoblot analysis.

Isotype lgG2b

Clonality Monoclonal

Clone Number 1H11-1H7

MW 78kDa


Uniprot ID P20029

NCBI NP_001156906.1

Entrez 14828

Dilution Range WB (1:2000), ICC/IF (1:100)


Expiration Date 12 months from date of receipt.




Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GRP78 Monoclonal Antibody, Clone 1H11-1H7. Tissue: Heat Shocked cervical cancer cells (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Endoplasmic reticulum lumen. Melanosome. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-GRP78 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.








Western Blot analysis of Human cell lysates showing detection of GRP78 protein using Mouse Anti-GRP78 Monoclonal Antibody, Clone 1H11-1H7. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody at 1:1000.



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GRP78 Monoclonal Antibody, Clone 1H11-1H7. Tissue: Heat Shocked cervical cancer cells (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Endoplasmic reticulum lumen. Melanosome. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-GRP78 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.



ELISA analysis using Mouse Anti-GRP78 Monoclonal Antibody, Clone 1H11-1H7. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody. Secondary Antibody: Goat anti-mouse IgG: HRP at 1:10000.