

Product Datasheet

Recombinant Human beta-Defensin 4 (rHuBD-4) (orb1494987)

Catalog Number	orb1494987
Category	Proteins
Description	<p>Defensins (alpha and beta) are cationic peptides with a broad spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The α-defensins are distinguished from the β-defensins by the pairing of their three disulfide bonds. To date, four human β-defensins have been identified; BD-1, BD-2, BD-3 and BD-4. The β-defensin proteins are expressed as the C-terminal portion of precursors and are released by proteolytic cleavage of a signal sequence. β-defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds. β-defensins are 3-5 kDa peptides ranging in size from 33-47 amino acid residues. BD-4 is expressed in testis, stomach, uterus, neutrophils, thyroid, lung and kidney. In addition to its direct antimicrobial activities, BD-4 is chemoattractant towards human blood monocytes.</p>
Form/Appearance	Lyophilized from a 0.2 μ m filtered concentrated solution in 20mM PB, pH 7.4, 130mM NaCl.
Buffer/Preservatives	Lyophilized from a 0.2 μ m filtered concentrated solution in 20mM PB, pH 7.4, 130mM NaCl.
Purity	>98% by SDS-PAGE and HPLC analyses.
Purification	>98% by SDS-PAGE and HPLC analyses.
Protein Sequence	EFELDRICGY GTARCRKKCR SQEYRIGRCP NTYACCLRKW DESLLNRTKP
MW	Approximately 6.0 KDa, a single non-glycosylated polypeptide chain containing 50 amino acids.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

Application notes	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at -20°C. Further dilutions should be made in appropriate buffered solutions.
Endotoxins	Less than 1EU/µg of rHuBD-4 as determined by LAL method.
Source	Escherichia coli.
Biological Activity	Determined by its ability to chemoattract human monocytes using a concentration range of 0.1-100.0 ng/ml, corresponding to a Specific Activity of 1 x 10 ⁴ IU/mg.
Storage	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.
Note	For research use only

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558