

Product Datasheet

NF-H Recombinant Rabbit Monoclonal Antibody (orb1499375)

Catalog Number orb1499375

Category Antibodies

Description NF-H Recombinant Rabbit Monoclonal Antibody

Target NEFH

Clonality Recombinant

Species/Host Rabbit

Isotype IgG

Conjugation Unconjugated

Reactivity Human, Mouse, Rat

Predicted Reactivity Human, Mouse, Rat

Form/Appearance Liquid

Concentration 1mg/ml

Buffer/Preservatives 0.01M TBS (pH7.4) with 1% rAlbumin, 0.02% Proclin300 and 50% Glycerol.

Purification Affinity purified by Protein A

Immunogen A synthesized peptide derived from human Neurofilament heavy polypeptide

(580-680aa)

UniProt ID P12036

MW 200 kDa





Tested applications IF, IHC-Fr, IHC-P, WB

Dilution range WB=1:500-2000, IHC-P=1:100-500, IHC-F=1:100-500, IF=1:100-500


Antibody Type Primary Antibody

Clone Number 4C3


Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only




Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-Neurofilament heavy polypeptide antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (orb1499375, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.





Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue using anti-Neurofilament heavy polypeptide antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (orb1499375, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.











Western blot analysis of Neurofilament heavy polypeptide on rat brain tissue lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (orb1499375, 1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:5000 dilution was used for 1 hour at room temperature.