

Product Datasheet

HSP40 Antibody: PerCP (orb151194)

Description Rabbit polyclonal to Hsp40 (PerCP). DnaJ/Hsp40 proteins have been preserved

throughout evolution and are important for protein translation, folding, unfolding, translocation, and degradation, primarily by stimulating the ATPase activity of chaperone proteins, Hsp70s. Because the ATP hydrolysis is essential for the activity of Hsp70s, DnaJ/Hsp40 proteins actually determine the activity of Hsp70s by stabilizing their interaction with substrate proteins. DnaJ/Hsp40 proteins all contain the J domain through which they bind to Hsp70s. Hsp40, also known as HDJ1, is a basic mammalian 40kDa heat shock protein which is not only homologous to the bacterial heat shock protein (DnaJ), but also yeast DnaJrelated proteins such as SCJ1, Sec63/Npl1, YDJ1 and SIS1 (2-5). Hsp 40 is inducible by stress including heat after which is moves from the cytoplasm to the nucleus and nucleoli; an intracellular pattern similar to Hsc70/Hsp70, the

mammalian homologues of the bacterial heat shock protein, DnaK..

Species/Host Rabbit

Reactivity Human

Conjugation PerCP

Tested Applications ELISA, ICC, IF, IHC, WB

Immunogen Recombinant purified full-length human HSP40 (no tags)

Target HSP40

Preservatives 95.64mM Phosphate, 2.48mM MES and 2mM EDTA

Storage Conjugated antibodies should be stored according to the product label

Note For research use only

Application notes 0.5 μg/ml of SPC-100 was sufficient for detection of HSP40 in 20 μg of heat

shocked HeLa cell lysate by colorimetric immunoblot analysis using Goat anti-

rabbit IgG:HRP as the secondary antibody.

Clonality Polyclonal

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>





MW 40kDa


Uniprot ID P25685

NCBI NP_006136.1

Entrez 3337

Dilution Range WB (1:2000), ICC/IF (1:100)

Expiration Date 12 months from date of receipt.




Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp40 Polyclonal Antibody. Tissue: Heat Shocked Cervical cancer cell line (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp40 Polyclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp40 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.



Western blot analysis of Human Cervical cancer cell line (HeLa) lysate showing detection of HSP40 protein using Rabbit Anti-HSP40 Polyclonal Antibody. Primary Antibody: Rabbit Anti-HSP40 Polyclonal Antibody at 1:1000.







Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp40 Polyclonal Antibody. Tissue: Heat Shocked Cervical cancer cell line (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp40 Polyclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: APC Goat Anti-Rabbit (red) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp40 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.

 $\begin{aligned} & \text{Email: } \underline{\text{info@biorbyt.com}}, \ \underline{\text{support@biorbyt.com}} \\ & \text{Phone: } \underline{+1 \ (415) \ 906\text{-}5211} \ \big| \ \text{Fax: } \underline{+1 \ (415) \ 651\text{-}8558} \end{aligned}$