

## Product Datasheet

### KDEL Antibody (PerCP) (orb151347)

<b>Catalog Number</b>	orb151347
<b>Category</b>	Antibodies
<b>Description</b>	<p>Rabbit polyclonal to KDEL (PerCP). The endoplasmic reticulum is part of a protein sorting pathway, or in essence, the transportation system of the eukaryotic cell. The majority of endoplasmic reticulum resident proteins are retained in the endoplasmic reticulum through a retention motif. This motif is composed of four amino acids at the C-terminal end of the protein sequence. The most common retention sequence is KDEL (lys-asp-glu-leu). Grp78 and Grp94 and PDI all share the C-terminal KDEL sequence. The presence of carboxy-terminal KDEL appears to be necessary for ER retention and appears to be sufficient to reduce the secretion of proteins from the ER..</p>
<b>Target</b>	KDEL
<b>Clonality</b>	Polyclonal
<b>Species/Host</b>	Rabbit
<b>Conjugation</b>	PerCP
<b>Reactivity</b>	Human, Mouse, Rat
<b>Concentration</b>	1 mg/ml
<b>Buffer/Preservatives</b>	95.46mM Phosphate, 2.48mM MES and 2mM EDTA
<b>Purification</b>	Protein A Purified
<b>Immunogen</b>	KDEL containing peptide immunogen
<b>Tested applications</b>	ICC, IF, IHC, WB
<b>Dilution range</b>	WB (1:1000), ICC/IF (1:100)

**Biorbyt Ltd.**

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

**Biorbyt LLC**

68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

**Application notes**

A 1:1000 dilution was sufficient for detection of KDEL-containing proteins in 20 µg of HeLa cell lysate by ECL immunoblot analysis using goat anti-mouse IgG as the secondary.

**Specificity**

Detects KDEL proteins, GRP94, Grp78, PDI and calreticulin. It may also see ERp57 and ERp29.

**Storage**


Conjugated antibodies should be stored according to the product label

**Note**


For research use only

**Expiration Date**

12 months from date of receipt.



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-KDEL Polyclonal Antibody. Tissue: Heat Shocked Cervical cancer cell line (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-KDEL Polyclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Endoplasmic reticulum. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-KDEL Antibody. (C) Composite. Heat Shocked at 42°C for 30 min.



1. PDI control antibody  
 2. KDEL control antibody clone 10C3  
 3. Grp78 control antibody  
 4. this product  
 5. Calreticulin control antibody  
 6. KDEL control antibody clone 10C3  
 7. Grp94 control antibody  
 Mixed human cell lysate (300ng/gel); 1/1000 dilutions;  
 KDEL(10C3) control antibody 1:500 dilution

Western blot analysis of Human Cell line lysates showing detection of KDEL protein using Rabbit Anti-KDEL Polyclonal Antibody. Primary Antibody: Rabbit Anti-KDEL Polyclonal Antibody at 1:1000, 1:500.

**Biorbyt Ltd.**


7 Signet Court, Swann Road  
 Cambridge  
 CB5 8LA  
 United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
 Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

**Biorbyt LLC**

68 TW Alexander Drive  
 Research Triangle Park  
 Durham  
 NC 27713  
 United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
 Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-KDEL Polyclonal Antibody. Tissue: Heat Shocked Cervical cancer cell line (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-KDEL Polyclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: R-PE Goat Anti-Rabbit (yellow) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Endoplasmic reticulum. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-KDEL Antibody. (C) Composite. Heat Shocked at 42°C for 30 min.

**Biorbyt Ltd.**

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

**Biorbyt LLC**

68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)