

Product Datasheet

GLI1 Rabbit Polyclonal Antibody (BF555) (orb1585214)

Catalog Number	orb1585214
Category	Antibodies
Description	GLI1 Rabbit Polyclonal Antibody (BF555) is a BF555 conjugated antibody targeting GLI1. This antibody is suitable for FC, IF. It exhibits reactivity with Human, Mouse, Rat samples.
Target	GLI1
Clonality	Polyclonal
Species/Host	Rabbit
Isotype	IgG
Conjugation	BF555
Reactivity	Human, Mouse, Rat
Predicted Reactivity	Bovine, Canine, Equine, Rabbit
Concentration	1mg/ml
Buffer/Preservatives	0.01M TBS (pH7.4) with 1% rAlbumin, 0.02% Proclin300 and 50% Glycerol.
Purification	Affinity purified by Protein A
Immunogen	KLH conjugated synthetic peptide derived from human GLI1/Zfp5 (601-700/1106aa)
Tested applications	FC, IF

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom


Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Dilution range	Flow-Cyt=1ug/Test, IF=1:100-500
Antibody Type	Primary Antibody
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Note	For research use only
Expiration Date	12 months from date of receipt.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30 min, Incubation: Anti-GLI1 Antibody, conjugated (orb1585214) at 1:100 overnight at 4°C, DAPI (5 ug/ml, blue) was used to stain the cell nuclei.



Paraformaldehyde-fixed, paraffin embedded (Mouse uterus), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30 min, Incubation: Anti-GLI1 Antibody, conjugated (orb1585214) at 1:100 overnight at 4°C, DAPI (5 ug/ml, blue) was used to stain the cell nuclei.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)