

Product Datasheet

Rabbit anti-Thrombomodulin Recombinant Monoclonal Antibody (orb1710821)

Catalog Number	orb1710821
Category	Antibodies
Description	Thrombomodulin Antibody
Target	Thrombomodulin
Clonality	Recombinant
Species/Host	Rabbit
Conjugation	Unconjugated
Reactivity	Mouse
Form/Appearance	Whole IgG
Concentration	500 µg/ml
Buffer/Preservatives	Borate Buffered Saline (BBS) pH 8.2 with 0.1% rAlbumin and 0.09% Sodium Azide
Immunogen	Residues 17-517 (ECD)
UniProt ID	P15306
Tested applications	FC, ICC, IHC, IP, WB
Dilution range	WB: 1:1,000, IP: 12 µl/1 mg lysate, IHC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections., ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections.
Application notes	Format: Whole IgG


Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)


Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Antibody Type	Primary Antibody
Clone Number	BLR260L
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Note	For research use only
Expiration Date	12 months from date of receipt.



Detection of mouse Thrombomodulin by western blot. Samples: Whole cell lysate (50 µg) from mIMCD-3, NIH 3T3, bEnd.3, C2C12, and CH27 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-Thrombomodulin recombinant monoclonal antibody (orb1710821) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG. Detection: Chemiluminescence with an exposure time of 3 minutes. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody.




Detection of mouse Thrombomodulin by immunohistochemistry. Sample: FFPE section of mouse gut. Antibody: Rabbit anti-Thrombomodulin recombinant monoclonal antibody (orb1710821). Secondary: HRP-conjugated goat anti-rabbit IgG.

Biorbyt Ltd.


7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44201223859353) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)

Biorbyt LLC.


68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+14159065211) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)



Detection of mouse Thrombomodulin (shaded) in 7OZ/3 cells by flow cytometry. Antibody: Rabbit anti-Thrombomodulin recombinant monoclonal antibody (orb1710821) or isotype control (unshaded). Secondary: DyLight® 650-conjugated goat anti-rabbit IgG.



Detection of mouse Thrombomodulin by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from bEnd.3 cells prepared using NETN lysis buffer. Antibodies: Rabbit anti-Thrombomodulin recombinant monoclonal antibody (orb1710821) used for IP at 12 µl/mg lysate.



Detection of mouse Thrombomodulin by immunocytochemistry. Sample: FFPE section of mouse 7OZ/3 cells. Antibody: Rabbit anti-Thrombomodulin recombinant monoclonal antibody (orb1710821). Secondary: HRP-conjugated goat anti-rabbit IgG.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)