

Product Datasheet

Recombinant Human Interleukin-2 receptor subunit alpha (IL2RA), partial (Active) (orb1785084)

Catalog Number	orb1785084
Category	Proteins
Description	This Recombinant Human Interleukin-2 receptor subunit alpha (IL2RA), partial (Active) spans the amino acid sequence from region 22-213aa. Purity: Greater than 95% as determined by SDS-PAGE.
Tag	C-terminal 10xHis-tagged
Form/Appearance	Lyophilized powder
Buffer/Preservatives	Lyophilized from a 0.2 µm filtered 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0
Purity	Greater than 95% as determined by SDS-PAGE.
Protein Sequence	ELCDDDPPEIPHATFKAMAYKEGTMLNCECKRGFRRIKSGSLYMLCTGNSSHSSWDNQCQCTSSATRNNTTKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPPWENEATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQPQLICTGEMETSQFPGEKPKQASPEGRPESETSC
Protein Length	Partial
UniProt ID	P01589
MW	23.6 kDa

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Application notes	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Endotoxins	Less than 1.0 EU/μg as determined by LAL method.
Source	Mammalian cell
Biological Origin	Homo sapiens (Human)
Biological Activity	① Measured by its binding ability in a functional ELISA. Immobilized Human IL2RA at 2 μg/mL can bind Anti-IL2RA recombinant antibody, the EC50 is 2.463-3.353 ng/mL. ② Measured by its binding ability in a functional ELISA. Immobilized Human IL2RA at 2 μg/mL can bind Human IL2, the EC50 is 1.693-2.039 ng/mL.
Expression Region	22-213aa
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Note	For research use only
Expiration Date	6 months from date of receipt.



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com


Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Measured by its binding ability in a functional ELISA. Immobilized Human IL2RA at 2 μ g/mL can bind Anti-IL2RA recombinant antibody, the EC50 is 2.463-3.353 ng/mL.



Measured by its binding ability in a functional ELISA. Immobilized Human IL2RA at 2 μ g/mL can bind Human IL2, the EC50 is 1.693-2.039 ng/mL.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558