

Product Datasheet

HA Tag Antibody (orb1787856)

Catalog Number	orb1787856
Category	Antibodies
Description	Purified Rabbit Polyclonal Antibody (Pab)
Target	KLH conjugated synthetic peptide encoding HA tag was used as antigen.
Clonality	Polyclonal
Species/Host	Rabbit
Isotype	Rabbit IgG
Conjugation	Unconjugated
Immunogen	KLH conjugated synthetic peptide encoding HA tag was used as antigen.
MW	45-50
Tested applications	WB
Dilution range	WB: 1:1000, WB: 1:1000, WB: 1:1000
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles
Note	For research use only
Expiration Date	12 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com


Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com


Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)



All lanes: Anti-HA Tag Antibody at 1:1000 dilution (upper) or GAPDH (lower). Lane 1: 293T/17 transfected with 12tag lysate (10ug). Lane 2: Non-transfected 293T/17 lysate (10ug). Lane 3: 12tag recombinant protein lysate (0. 1ug). Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 45-50 kDa. Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes: Anti-HA Tag Antibody at 1:1000 dilution (upper) or GAPDH (lower). Lane 1: 293T/17 transfected with 12tag lysate (10ug). Lane 2: Non-transfected 293T/17 lysate (10ug). Lane 3: 12tag recombinant protein lysate (0. 1ug). Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 45-50 kDa. Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes: Anti-HA Tag Antibody at 1:1000 dilution (upper) or GAPDH (lower). Lane 1: 293T/17 transfected with 12tag lysate (0. 5ug). Lane 2: Non-transfected 293T/17 lysate (0. 5ug). Lane 3: 12tag recombinant protein lysate (0. 1ug). Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 45-50 kDa. Blocking/Dilution buffer: 5% NFDm/TBST.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558