

Product Datasheet

RANBP9 Antibody (orb1925554)

Catalog Number	orb1925554
Category	Antibodies
Description	RANBP9 Antibody
Clonality	Monoclonal
Species/Host	Mouse
Isotype	IgG1,k
Conjugation	Unconjugated
Reactivity	Human, Mouse
Form/Appearance	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
UniProt ID	Q96S59
MW	77847 Da
Tested applications	IHC-P, WB
Dilution range	IHC-P - 1:100-500, WB - 1:2000-1:4000
Specificity	This RANBP9 antibody is generated from a mouse immunized with a recombinant protein between 1-388 amino acids from the human RANBP9.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles
Note	For research use only

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States


Email: info@biorbyt.com, support@biorbyt.com

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Expiration Date 12 months from date of receipt.



All lanes: Anti-RANBP9 Antibody at 1:2000-1:4000 dilution. Lane 1: HeLa whole cell lysate. Lane 2: U-2OS whole cell lysate. Lane 3: Jurkat whole cell lysate. Lane 4: CCRF-CEM whole cell lysate. Lane 5: MDA-MB-231 whole cell lysate. Lane 6: MOLT-4 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 78 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



Staining RANBP9 in human heart tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)