

Product Datasheet

HNRNPA1L2 Antibody (N-Term) (orb1926199)

Catalog Number	orb1926199
Category	Antibodies
Description	Purified Rabbit Polyclonal Antibody (Pab)
Target	HNRNPA1L2
Clonality	Polyclonal
Species/Host	Rabbit
Isotype	Rabbit IgG
Conjugation	Unconjugated
Reactivity	Human
Form/Appearance	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Immunogen	This HNRNPA1L2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 22-56 amino acids from human HNRNPA1L2.
UniProt ID	Q32P51
MW	34225 Da
Tested applications	IHC-P, WB
Dilution range	WB - 1:2000, IHC-P - 1:25

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States


Email: info@biorbyt.com, support@biorbyt.com

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

- Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles
- Note** For research use only
- Expiration Date** 12 months from date of receipt.



All lanes: Anti-HNRNPA1L2 Antibody (N-Term) at 1:2000 dilution. Lane 1: HeLa whole cell lysate. Lane 2: Jurkat whole cell lysate. Lane 3: MCF-7 whole cell lysate. Lane 4: HepG2 whole cell lysate. Lane 5: K562 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 34 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



Staining HNRNPA1L2 in human breast carcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558