

Product Datasheet

LGALS3BP Antibody (Center) (orb1932111)

Catalog Number	orb1932111
Category	Antibodies
Description	Affinity Purified Rabbit Polyclonal Antibody (Pab)
Target	LGALS3BP
Clonality	Polyclonal
Species/Host	Rabbit
Isotype	Rabbit IgG
Conjugation	Unconjugated
Reactivity	Human, Mouse
Form/Appearance	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Immunogen	This LGALS3BP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 383-412 amino acids from the Central region of human LGALS3BP. Antigen Region: 383-412 aa.
UniProt ID	Q08380
MW	65331 Da
Tested applications	IHC-P, WB
Dilution range	WB - 1:2000, IHC-P-Leica - 1:100
Antibody Type	Primary Antibody

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom


Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles
Note	For research use only
NCBI	NP_005558.1
Expiration Date	12 months from date of receipt.



Immunohistochemical analysis on paraffin-embedded Human prostate tissue was performed on the Leica BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody (1:100) for 15 min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



All lanes: Anti-LGALS3BP Antibody (Center) at 1:2000 dilution.
Lane 1: HeLa whole cell lysate. Lane 2: Human plasma lysate. Lane 3: Mouse lung lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 65 kDa.
Blocking/Dilution buffer: 5% NFDM/TBST.

Biorbyt Ltd.


7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



All lanes: Anti-LGALS3BP Antibody (Center) at 1:2000 dilution.
Lane 1: Human liver tissue lysate. Lane 2: Mouse lung tissue lysate. Lane 3: Human plasma tissue lysate. Lane 4: NIH/3T3 whole cell lysate. Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 65 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)