

Product Datasheet

Rabbit Bovine IgM (Fc specific), conjugated with FITC Antibody (orb20670)

Catalog Number orb20670

Category Antibodies

Description Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal rabbit antiserum

to Bovine IgM, Fc specific. In immunocytochemical and immunohistochemical staining for the detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgM antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of bovine origin known to be of the IgM isotype in the middle layer of the indirect test procedure. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually

between 1:20 and 1:80.

Clonality Polyclonal

Species/Host Rabbit

Conjugation FITC

Reactivity Bovine

Form/Appearance Fluorescein-coupled purified hyperimmune rabbit IgG lyophilized from a solution

in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

RRID AB_10931198

Antibody Type Secondary Antibody

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: $\underline{info@biorbyt.com}$, $\underline{support@biorbyt.com}$ Phone: $\underline{+44~(0)1223~859353}$ | Fax: $\underline{+1~(415)~651}$ -8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Biorbyt.com

Source

Purified normal IgM isolated from pooled bovine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at $+4^{\circ}$ C; prolonged storage at or below -20° C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note

For research use only

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

$$\begin{split} & \text{Email: } \underline{\text{info@biorbyt.com}}, \underline{\text{support@biorbyt.com}} \\ & \text{Phone: } \underline{+44~(0)1223~859353} \mid \text{Fax: } \underline{+1~(415)~651-8558} \end{split}$$

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558