

## **Product Datasheet**

## Goat anti Rabbit IgG (heavy and light chains), conjugated with Horseradish peroxidase (orb21461)

## Description

Horseradish peroxidase-conjugated IgG fraction of polyclonal goat antiserum to Rabbit IgG, heavy and light chains. In enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG, antigen or antibody, of appropriately treated cell and tissue substrates at the cellular and subcellular level. In non-isotopic assay methodology (e.g. ELISA) to identify and measure a specific IgG in rabbit serum or other body fluid. In electron microscopy, since the complex between the conjugated antibody and the antigen also has electrondense properties. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:500; in ELISA between 1:5,000 and 1:15,000; in blotting up to 1:200,000.

Species/Host Goat

**Reactivity** Rabbit

**Conjugation** HRP

**Tested Applications** DOT, ELISA, ICC, IHC, WB

**Preservatives** Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

**Form/Appearance** Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



## **Biorbyt.com**

**Storage** 

The lyophilised conjugate is shipped at ambient temperature and may be stored at  $+4^{\circ}$ C; prolonged storage at or below  $-20^{\circ}$ C. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at  $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the

performance of the product.

**Note** For research use only

**Clonality** Polyclonal

Antibody Type Secondary Antibody

**Source** Purified normal IgG isolated from pooled rabbit serum. Freund's complete

adjuvant is used in the first step of the immunization procedure.

**Hazard Information** This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

**Expiration Date** 12 months from date of receipt.