

Product Datasheet

Rabbit anti Goat IgG (heavy and light chains) (non Human and Mouse), conjugated with FITC (orb21499)

Catalog Number orb21499

Description Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal rabbit antiserum

to goat IgG, heavy and light chains, adsorbed for cross-reactivity with human and mouse immunoglobulins. To identify and measure IgG, antigen or antibody,

at the cellular and subcellular level by immuno- fluorescence staining of

appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of goat origin in the middle layer of the indirect test procedure. This immunoconjugate is not

pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be

avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually between 1:20 and 1:80.

Species/Host Rabbit

Reactivity Goat

Conjugation FITC

Preservatives FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added

Form/Appearance FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added



Biorbyt.com

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at $+4^{\circ}$ C; prolonged storage at or below -20° C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note For research use only

Clonality Polyclonal

Antibody Type Secondary Antibody

Source Purified normal IgG isolated from pooled goat serum. Freund's complete

adjuvant is used in the first step of the immunization procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.