

Product Datasheet

Rabbit anti Sheep IgM (Fc specific), conjugated with FITC (orb21558)

Catalog Number orb21558

Description Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal rabbit antiserum

to sheep IgM, Fc specific. In immunocytochemical and immunohistochemical staining for the detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgM antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of sheep origin known to be of the IgM isotype in the middle layer of the indirect test procedure. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually

between 1:20 and 1:80.

Species/Host Rabbit

Reactivity Sheep

Conjugation FITC

Preservatives Fluorescein-coupled purified hyperimmune rabbit IgG lyophilized from a solution

in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Form/Appearance Fluorescein-coupled purified hyperimmune rabbit IgG lyophilized from a solution

in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Biorbyt.com

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note For research use only

Clonality Polyclonal

Antibody Type Secondary Antibody

Source Purified normal IgM isolated from pooled sheep serum. Freund's complete

adjuvant is used in the first step of the immunization procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.