

Product Datasheet

Goat anti Mouse IgG1 IgG2a IgG2b IgG3 IgA IgM (Fc specific), conjugated with Biotin (orb21642)

Catalog Number orb21642

Description Biotin-conjugated IgG fraction of polyclonal goat antiserum to Mouse IgG, IgA and IgM, Fc specific. In immunocytochemical and immunohistochemical staining of immunoglobulins at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of mouse origin in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure immunoglobulins in mouse serum or other body fluids. The absence of activity to the common Ig/Fab subunit prevents the reaction of this conjugate with immunoglobulins binds to Fc receptors on non-lymphoid cells. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:500; in ELISA and comparable non-precipitating antibody-binding assays between 1:1,000 and 1:7,000.

Species/Host Goat

Reactivity Mouse

Conjugation Biotin

Preservatives Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Form/Appearance	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio
Storage	The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.
Note	For research use only
Clonality	Polyclonal
Antibody Type	Secondary Antibody
Source	Purified normal IgG an homogenous IgA and IgM isolated from mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Hazard Information	This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no liability for any inaccuracies or omissions in this information.
Expiration Date	12 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)