

Product Datasheet

Rabbit anti Monkey IgG Fab, conjugated with FITC (orb21650)

Catalog Number orb21650

Description Fluorescein isothiocyanate-conjugated F(ab')2 fragments IgG antibodies of

polyclonal goat antiserum to human Fab IgG. Direct staining of fixed cell and tissue substrates, to demonstrate the intracellular presence of free of Ig bound subunits of both kappa and lambda type. The absence of the Fc domain in the conjugate ensures minimal interaction with the tissue components and cell surfaces other than the primary antibody activity. This conjugate is primarily intended for use in cell surface membrane staining procedures, to identify and quantitate Ig on B cells, especially if interference by Fc activity is expected. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually

between 1:20 and 1:80.

Species/Host Rabbit

Reactivity Monkey

Conjugation FITC

Preservatives FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added as it may

interfere with he antibody activity.

Form/Appearance FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added as it may

interfere with he antibody activity.



Biorbyt.com

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note For research use only

Clonality Polyclonal

Antibody Type Secondary Antibody

Source Purified normal Fab of monkey polyclonal IgG. Freund's complete adjuvant is

used in the first step of the immunization procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.

Phone: $\pm 1 (415) 906-5211$ | Fax: $\pm 1 (415) 651-8558$