

Product Datasheet

Rabbit Goat IgG (Fc specific), conjugated with Horseradish peroxidase Antibody (orb21717)

Catalog Number orb21717

Category Antibodies

Description Horseradish peroxidase-conjugated IgG fraction of polyclonal rabbit antiserum to

goat IgG, Fc specific. In enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgG antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of goat origin known to be of the IgG isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgG in goat serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:250; in ELISA and comparable non-precipitating antibody-binding assays between

1:2,000 and 1:8,000.

Clonality Polyclonal

Species/Host Rabbit

Conjugation HRP

Reactivity Goat

Form/Appearance Horseradish peroxidase-coupled purified hyperimmune rabbit IgG lyophilized

from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative

added, as it may interfere with the antibody activity.

RRID AB_10937496

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: $\underline{info@biorbyt.com}$, $\underline{support@biorbyt.com}$ Phone: $\underline{+44~(0)1223~859353}$ | Fax: $\underline{+1~(415)~651}$ -8558

Biorbyt LLC

United States

68 TW Alexander Drive Research Triangle Park Durham NC 27713

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558





Antibody Type Secondary Antibody

Source Purified normal IgG isolated from pooled goat serum. Freund's complete

adjuvant is used in the first step of the immunization procedure.

Storage The lyophilized conjugate is shipped at ambient temperature and may be stored

at $+4^{\circ}$ C; prolonged storage at or below -20° C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect

the performance of the immunoconjugate.

Note For research use only

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558