

Product Datasheet

Goat anti Human IgE (Fc specific), conjugated with FITC (orb21834)

Catalog Number orb21834

Description Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal goat antiserum

to Human IgE Fc specific. In immunocytochemical and immunohistochemical staining for the detection of IgE at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to identify and measure IgE in human serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually between 1:20 and 1:80, depending on the method

used.

Species/Host Goat

Reactivity Human

Conjugation FITC

Preservatives FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Form/Appearance FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Storage The lyophilized conjugate is shipped at ambient temperature and may be stored

at $+4^{\circ}$ C; prolonged storage at or below -20° C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect

the performance of the immunoconjugate.





Note For research use only

Clonality Polyclonal

Antibody Type Secondary Antibody

Source Pools of purified homogenous IgE isolated from human serum. Freund's

complete adjuvant is used in the first step of the immunization procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>