

Product Datasheet

Goat Human IgG Fab, conjugated with TRITC Antibody (orb21866)

Catalog Number orb21866

Category Antibodies

Description Tetramethylrhodamine isothiocyanate-conjugated IgG fraction of polyclonal goat

antiserum to human Fab of IgG. Direct staining of fixed cell and tissue substrates; to demonstrate the intracellular presence of free or Ig-bound subunits of both kappa or lambda type. In general this conjugate is not recommended as direct or indirect screening reagent for If isotypes on surface

membranes of vital lymphoid cells. The activity to the common Ig/Fab subunit may result in the staining of immunoglobulins bound to the Fc-receptors on non-lymphoid cells. Combinations of isotype-specific reagents should be used instead for this purpose. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are

usually between 1:10 and 1:40.

Clonality Polyclonal

Species/Host Goat

Conjugation TRITC

Reactivity Human

Form/Appearance TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added as it may

interfere with he antibody activity.

RRID AB_10949048

Antibody Type Secondary Antibody





Source

Purified Fab of normal IgG isolated form human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note

For research use only

Expiration Date

12 months from date of receipt.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558