

Product Datasheet

Goat anti Guinea Pig IgG (heavy and light chains), conjugated with FITC (orb22037)

Description

Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal goat antiserum to guinea pig IgG, heavy and light chains. To identify and measure IgG, antigen or antibody, at the cellular and subcellular level by immuno- fluorescence staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of guinea pig origin in the middle layer of the indirect test procedure. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually between 1:20 and 1:120.

Species/Host Goat

Conjugation FITC

Preservatives FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added.

Form/Appearance FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added.

Storage The lyophilized conjugate is shipped at ambient temperature and may be stored

at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH

7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It

will not affect the performance of the immunoconjugate.

Note For research use only

Clonality Polyclonal





Antibody Type Secondary Antibody

Source Purified normal IgG isolated from pooled guinea pig serum. Freund's complete

adjuvant is used in the first step of the immunization procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

......

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.