

Product Datasheet

Goat IgG anti Hamster IgG (heavy and light chains), conjugated with Horseradish peroxidase (orb22039)

Description	Horseradish peroxidase-conjugated IgG fraction of polyclonal goat antiserum to hamster IgG, heavy and light chains. In enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG, antigen or antibody, of appropriately treated cell and tissue substrates at the cellular and subcellular level. In non-isotopic assay methodology (e.g. ELISA) to identify and measure a specific IgG in hamster serum or other body fluid. In electron microscopy, since the complex between the conjugated antibody and the antigen also has electron-dense properties. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1: 500; in ELISA and comparable non-precipitating antibody-binding assays between 1:1,000 and 1:5,000.
Species/Host	Goat
Reactivity	Hamster
Conjugation	HRP
Tested Applications	DOT, ELISA, ICC, WB
Preservatives	Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Form/Appearance	Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Storage	The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.
Note	For research use only
Clonality	Polyclonal
Antibody Type	Secondary Antibody
Source	Goats were immunized with purified normal IgG isolated from pooled hamster serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Hazard Information	This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no liability for any inaccuracies or omissions in this information.
Expiration Date	12 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)