

Product Datasheet

Goat anti Mouse IgG1 IgG2a IgG2b IgG3 IgA IgM (heavy and light chains) (non Human), conjugated with Biotin (orb22084)

Catalog Number orb22084

DescriptionBiotin-conjugated IgG fraction of polyclonal goat antiserum to Mouse IgG, IgA

and IgM, heavy and light chains, adsorbed for human cross-reactivity. In immunocytochemical and immunohistochemical staining of monoclonal mouse monoclonal antibodies in the presence of human cell or serum components at the cellular and subcellular level. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not prediluted. The optimum working dilution of each conjugate should be established

by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions are usually between 1:100 and 1:500; in ELISA and comparable non-precipitating antibody-binding assays between 1:1,000 and

1:10,000.

Species/Host Goat

Reactivity Mouse

Conjugation Biotin

PreservativesBiotin-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Form/Appearance Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.



Biorbyt.com

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note For research use only

Clonality Polyclonal

Antibody Type Secondary Antibody

Source Purified polyclonal mouse IgG, and homogenous IgA and IgM isolated from

mouse serum. Freund's complete adjuvant is used in the first step of the

immunization procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.