

Product Datasheet

Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (heavy and light chains) (non Human and Mouse), conjugated with TRITC (orb22096)

Catalog Number orb22096

Description Tetramethylrhodamine isothiocyanate-conjugated IgG fraction of polyclonal goat

antiserum to rat IgG, IgA and IgM, heavy and light chains, adsorbed for human and mouse cross-reactivity. Direct immunofluorescence staining of cytoplasmic

Ig of appropriately treated cell and tissue substrates; to demonstrate

immunoglobulins or specific antibodies in cells and tissues; to identify circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of rat origin in the middle layer of the indirect test procedure. This immunoconjugate is not prediluted. The optimum working dilution of each conjugate should be established

by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the

specific signal. Working dilutions are usually between 1:10 and 1:40.

Species/Host Goat

Reactivity Rat

Conjugation TRITC

Preservatives TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Form/Appearance TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.



Biorbyt.com

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at $+4^{\circ}$ C; prolonged storage at or below -20° C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

Note For research use only

Clonality Polyclonal

Antibody Type Secondary Antibody

Source Purified polyclonal rat IgG, and homogenous IgA and IgM isolated from rat

serum. Freund's complete adjuvant is used in the first step of the immunization

procedure.

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>