

Product Datasheet

Multi-rAb HRP-Goat Rabbit Recombinant Secondary Antibody (H+L) (orb2302687)

Catalog Number	orb2302687
Category	Antibodies
Description	This Multi-rAb antibody targets to Rabbit IgG (H+L) in ELISA and Western blot. It is highly specific to rabbit IgG and shows minimal reactivity for mouse, rat, human, pig, and bovine IgGs.
Clonality	Recombinant
Species/Host	Goat
Conjugation	HRP
Reactivity	Rabbit
Form/Appearance	Liquid
Concentration	1.0 mg/mL
Buffer/Preservatives	PBS with 50% glycerol, 10 mg/mL rAlbumin, 0.1% Proclin-300, pH 7.4.
Purity	The antibody was purified from culture media supernatant by immunoaffinity chromatography using Protein G beads.
Tested applications	DOT, ELISA, WB
Dilution range	1:10,000-1:100,000 for ELISA, 1:10,000-1:200,000 for western blotting with ECL substrates (1:10,000-1:40,000 is suggested for most system).
Antibody Type	Recombinant Antibody
Storage	Store at -20°C. Stable for one year after shipment.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom


Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)


Note For research use only



Various lysates were subjected to SDS-PAGE followed by western blot with rabbit anti-TOM70 polyclonal antibody at dilution of 1:10000. Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687) was used at 1:20000 for detection.



Various lysates were subjected to SDS-PAGE followed by western blot with rabbit anti-PSMB1 polyclonal antibody at dilution of 1:10000. Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687) was used at 1:20000 for detection.



Various lysates were subjected to SDS-PAGE followed by western blot with rabbit anti-Beta Catenin polyclonal antibody at dilution of 1:50000. Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687) was used at 1:20000 for detection.

Biorbyt Ltd.


7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States


Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Various lysates were subjected to SDS-PAGE followed by western blot with rabbit anti-TDP43 recombinant antibody at dilution of 1:20000. Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687) was used at 1:20000 for detection.



PC-3 and Calyculin A treated PC-3 cell lysates were subjected to SDS-PAGE followed by western blot with rabbit anti-Phospho-EIF4B (Ser406) Recombinant antibody at dilution of 1:20000. Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687), leading competitor's non cross-adsorbed and cross-adsorbed secondary antibodies were all used at 0.05 µg/mL for detection.



HEK-293 and HepG2 cell lysates were subjected to SDS-PAGE followed by western blot with rabbit anti-ZO-1 polyclonal antibody at dilution of 1:50000. Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687), leading competitor's non cross-adsorbed and cross-adsorbed secondary antibodies were all used at 0.05 µg/mL for detection.

Biorbyt Ltd.


7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558


Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Various lysates were subjected to SDS-PAGE followed by western blot with Beta tubulin rabbit recombinant antibody at dilution of 1:20000. Two batches of Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687) were used at 1:20000 for detection.



Rabbit total IgG, Mouse total IgG, Rat total IgG, Pig total IgG, Human total IgG, Bovine total IgG were coated at 100 ng/well. 0.125 µg/mL Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (orb2302687) was used for detection. The result indicates that orb2302694 is highly specific for rabbit IgG and does not react with other species tested in the experiment.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)