

## **Product Datasheet**

# Recombinant S100B (Astrocyte and Melanoma Marker) Antibody (orb2317128)

Catalog Number orb2317128

**Category** Antibodies

**Description** Recombinant S100B (Astrocyte and Melanoma Marker) Antibody

Target S100B

**Clonality** Monoclonal

Species/Host Rabbit

**Isotype** IgG

**Conjugation** Unconjugated

**Reactivity** Bovine, Canine, Human, Mouse, Rat

**Form/Appearance** 200ug/ml of Ab produced in HEK293 cell mammalian-based expression system.

Prepared in 10mM PBS with 0.05% rAlbumin & 0.05% azide. Also available

WITHOUT rAlbumin & azide at 1.0mg/ml.

**Concentration** Purified Ab with BSA and Azide at 200ug/ml

Immunogen Recombinant full-length human S100B protein

UniProt ID P04271

**MW** 10-12kDa

**Tested applications** IHC, WB

#### **Biorbyt Ltd.**

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

#### **Biorbyt LLC**

68 TW Alexander Drive Research Triangle Park Durham

NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



### **Biorbyt.com**

**Dilution range** Western Blot (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for

30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be

determined.

**Application notes** Positive Control: Human brain or melanoma

**Specificity** S100 belongs to the family of calcium binding proteins. S100A and S100B

proteins are two members of the S100 family. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epitope located on the beta-chain (i.e. in S-100A and S-100B) but not on the alpha-chain of S-100 (i.e. in S-100A and S100A0). This antibody can be used to localize S-100A and S-100B in various tissue sections. S-

100 protein has been found in normal melanocytes, Langerhans cells,

histiocytes, chondrocytes, lipocytes, skeletal and cardiac muscle, Schwann cells, epithelial and myoepithelial cells of the breast, salivary and sweat glands, as well as in glial cells. Neoplasms derived from these cells also express S-100 protein, albeit non-uniformly. A large number of well-differentiated tumors of the

salivary gland, adipose and cartilaginous tissue, and Schwann cell-derived tumors express S-100 protein. Almost all malignant melanomas and cases of

histiocytosis X are positive for S-100 protein.

**Antibody Type** Recombinant Antibody

Clone Number S100B/1706R

**Expression System** HEK293 cells

**Storage** Antibody with azide store at 2 to 8 °C. Antibody without azide store at -20 to -80

°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

**Note** For research use only

Entrez 6285

**Biorbyt Ltd.** 

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

$$\begin{split} & \text{Email: } \underline{\text{info@biorbyt.com}}, \underline{\text{support@biorbyt.com}} \\ & \text{Phone: } \underline{+44\ (0)1223\ 859353} \mid \text{Fax: } \underline{+1\ (415)\ 651\text{-}8558} \end{split}$$

**Biorbyt LLC** 

68 TW Alexander Drive Research Triangle Park Durham NC 27713

NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558