

## **Product Datasheet**

## **Anti-His tag (C-term) [3D5] (orb256371)**

**Description** Mouse monoclonal antibody to His tag

**Species/Host** Mouse

**Conjugation** Unconjugated

**Tested Applications** ELISA, FACS, FC, WB

**Immunogen** Purified his tagged Fab fragment.

**Target** His tag (C-term)

**Preservatives** PBS with 0.02% Proclin 300.

**Concentration** 1 mg/ml

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

**Note** For research use only

**Isotype** IgG2a

**Clonality** Monoclonal

Clone Number 3D5


**Purity** Purified

**Expiration Date** 12 months from date of receipt.


Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558








Immunofluorescence staining of HEK293 cells with Anti-His tag (C-term) 3D5. Immunofluorescence analysis of paraformaldehyde fixed HEK293 cells transfected with His tag expressing plasmid (A) and non-transfected HEK293 cells (B), permeabilized with 0.15% Triton stained with the chimeric rabbit IgG version of 3D5 (orb256373) (1:200 dilution) for 1h followed by Alexa Fluor® 488 secondary antibody (1:1000 dilution), showing cytoplasmic staining. The nuclear stain is DAPI (blue).



orb256366 (1  $\mu$ g/ml, right panel) staining of Multitag protein at 25 ng (A), 10 ng (B) and 5 ng (C) amounts. Primary incubation was 1 hour. Detected by chemiluminescence.



Western Blot using Anti-His tag (C-term) antibody 3D5. His tag expressing plasmid transfected (A) and non-transfected (B) HEK293 cells lysate (3g protein in RIPA buffer) were resolved on a SDS PAGE gel and blots were probed with the chimeric rabbit version of 3D5 (orb256373) at 0.01  $\mu$ g/ml before detection using an anti-rabbit secondary antibody. A primary incubation of 1h was used and protein was detected by chemiluminescence.