

Product Datasheet

RXR gamma Antibody / RXRG (orb2634839)

Catalog Number orb2634839

Description Two families of retinoid receptors, RARs and RXRs, have been identified. Retinoic

acid receptors (RARs) include RARa, RARb and RARg, each of which have a high affinity for all trans-retinoic acids and belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D3 receptor and ecdysone receptor. The ligand-binding domains of the RARs are highly conserved and RAR isoforms are expressed in distinct patterns throughout development and in the mature organism. Members of the retinoid X receptor (RXR) family, RXR, are activated by 9-cis-RA, a stereo- and photo-isomer of all trans-RA that is expressed in vivo in both liver and kidney and may represent a widely used hormone. As is true for the RAR subfamily, the RXR receptors are closely related to each other both in their DNA-binding and ligand-binding domains and are

encoded by separate genes at distinct chromosomal loci.

Species/Host Mouse

Reactivity Human

Conjugation Unconjugated

Tested Applications ELISA, FACS, IF

Immunogen Recombinant full-length human RXRG protein was used as the immunogen for

the RXR gamma antibody.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Application notes Optimal dilution of the RXR gamma antibody should be determined by the

researcher.

Formula 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide

Isotype Mouse IgG1





Clonality Monoclonal

Clone Number PCRP-RXRG-5H4

Antibody Type Primary Antibody


Uniprot ID P48443

Hazard Information This RXR gamma antibody is available for research use only.


Dilution Range ELISA (order BSA-free format for coating), Flow cytometry: 1-2ug/million

cells,Immunofluorescence: 1-2ug/ml

Expiration Date 12 months from date of receipt.



Immunofluorescent staining of PFA-fixed human HeLa cells using RXR gamma antibody (green, clone PCRP-RXRG-5H4) and phalloidin (red).




FACS staining of PFA-fixed human HeLa cells with RXR gamma antibody (blue, clone PCRP-RXRG-5H4) and isotype control (red).





Human Protein Microarray Specificity Validation



Analysis of HuProt (TM) microarray containing more than 19000 full-length human proteins using RXR gamma antibody (clone PCRP-RXRG-5H4). These results demonstrate the foremost specificity of the PCRP-RXRG-5H4 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt (TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt (TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free RXR gamma antibody (clone PCRP-RXRG-5H4) as confirmation of integrity and purity.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>