

Product Datasheet

Interleukin 15 Antibody / IL15 (orb2635055)

Catalog Number orb2635055

Description Interleukin-15 (IL-15), also designated IL-T, is a cloned cytokine which shares

several biological activities but no sequence homology with IL-2. Human, mouse and simian IL-15 cDNA clones have been isolated and characterized. All 3

species encode a 162 amino acid residue precursor protein containing a 48 amino acid leader that is cleaved to generate the mature form of IL-15. IL-15 stimulates the proliferation of T cells and NK cells, while enhancing B cell

expansion and antibody production. Unlike IL-2, IL-15 is not produced by lymphocytes, but appears to be produced by macrophages, epithelial lines, muscle and placenta. IL-15 has also been shown to be a chemoattractant for

human blood T lymphocytes and to be able to induce lymphokine-activated killer (LAK) activity in NK cells as well as to be able to induce the generation of

cytolytic effector cells. Studies have shown that IL-15 is the only other cytokine

that shares the subunit.

Species/Host Mouse

Reactivity Human

Conjugation Unconjugated

Tested Applications IHC-P

Immunogen A recombinant protein fragment was used as the immunogen for the Interleukin

15 antibody.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Application notes Optimal dilution of the Interleukin 15 antibody should be determined by the

researcher.

Formula 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide





Isotype Mouse IgG1, kappa

Clonality Monoclonal

Clone Number IL15/4353


Antibody Type Primary Antibody

Uniprot ID P40933


Hazard Information This Interleukin 15 antibody is available for research use only.

Dilution Range Immunohistochemistry (FFPE): 1-2ug/ml

Expiration Date 12 months from date of receipt.



SDS-PAGE analysis of purified, BSA-free Interleukin 15 antibody (clone IL15/4353) as confirmation of integrity and purity.




IHC staining of FFPE human adrenal gland tissue with Interleukin 15 antibody (clone IL15/4353). HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>





Human Protein Microarray Specificity Validation



Analysis of HuProt (TM) microarray containing more than 19000 full-length human proteins using Interleukin 15 antibody (clone IL15/4353). These results demonstrate the foremost specificity of the IL15/4353 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt (TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt (TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>