

Product Datasheet

One step Polymer HRP Goat IgG (H+L) (orb2652844)

Description IBSC One-Step rabbit anti-Goat IgG (H+L); biotin/avidin free system stains

membranes, cytoplasmic and nuclear antigens. It provides the user with a rapid

and easy to use IHC detection system.

Tested Applications ICC, IHC

Storage 2-8 °C, Do Not Freeze

Note For research use only

Application notes

Procedure: IHC/ICC procedure for frozen, paraffin sections and cell smears. 1. Deparafinize and hydrate tissue sections through xylene or other clearing agents and graded alcohols. (For frozen sections or cell smears; use unfixed, acetone fixed or appropriate fixative for the antigen in question; for cell smears it may be necessary to permealize the cell by detergent, please refer to antibody protocol). 2. Wash 2-3 with distilled or deionized water. 3. Incubate sections/cell smear with Endoblocker (Not supplied) for 5-10 minutes at room temperature (RT). Wash with distilled water 3X. 4. Note: If antigen retriever (Trypsin AR-6541, Pronase AR-6542, Pepsin AR-6543, Citrate buffer AR-6544, Buffer w EDTA pH 8.5 AR-6545, Tris buffer pH 10 AR-6546) is required it can be applied at this step. Please refer to data sheet for the primary antibody. 5. Wash slide with PBS or Tris saline buffer (with 0.02-0.05% nonionic detergent, Triton X100, Tween 20 or NP-40) or washing buffer (Immuno Automation buffer IBSC cat # AR-6561) 3X. 6. Incubate sections/ cell smear in Protein blocking solution (Not supplied), for 5-10 minutes at RT. 7. Incubate sections/cell smear with primary antibody (NOT SUPPLIED) for 20-30 minutes at 8. Wash slide with PBS 5-7X 9. Incubate with One-Step HRP polymer (Supplied) for 20-30 minutes at RT. 10. Wash slide 5-7 times with buffer. Caution: Peroxidase reagents are destroyed by sodium azide and should be avoided in all buffers and regents. 11. Wash slide with deionized or distilled for 2-3X. 12. Incubate with AEC or DAB chromogen reagent (Not supplied). 13. Wash 5-7X with buffer. 14. Incubate with counterstain (Not supplied). 15. Wash slide with tap water distilled water. 16. Mount slide with appropriate mounting medium.

Expiration Date

12 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: +44 (0) 1223 859-353 | Fax: Fax: +1 (415) 651-8558

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558