

Product Datasheet

TAK1 (Phospho-T184) Antibody (orb315597)

Catalog Number orb315597

Category Antibodies

Description The TAK1 (Phospho-T184) Antibody is suitable for IF, IHC, WB. It is a Polyclonal,

Unconjugated antibody which raised against KLH-conjugated synthetic

phosphopeptide corresponding to residues surrounding T184 of human TAK1 protein. The exact sequence is proprietary. Purification: The antibody was

purified by immunogen affinity chromatography.

Clonality Polyclonal

Species/Host Rabbit

Conjugation Unconjugated

Reactivity Bovine, Human, Mouse, Rat

Form/Appearance Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.01% sodium azide.

Purification The antibody was purified by immunogen affinity chromatography.

Immunogen KLH-conjugated synthetic phosphopeptide corresponding to residues

surrounding T184 of human TAK1 protein. The exact sequence is proprietary.

UniProt ID 043318, Q62073, P0C8E4

Tested applications IF, IHC, WB

Dilution range WB: 1-500:1000, IHC-P: 1-100:200

Antibody Type Primary Antibody

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.







Note For research use only

Entrez 6885, 26409, 313121, 100910771


Expiration Date 12 months from date of receipt.



Western blot analysis of TAK1 (Phospho-T184) expression in mouse muscle (A) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 70 kD)



Immunohistochemical analysis of TAK1 (Phospho-T184) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (Phospho-H 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.




Immunofluorescent analysis of TAK1 (Phospho-T184) staining in LS8 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>







Direct ELISA antibody dose-response curve using Anti-TAK1 (Phospho-T184) Antibody. Antigen (Phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>