

Product Datasheet

Mouse IgG (H&L) Antibody Peroxidase Conjugated (orb346678)

Description Mouse IgG (H&L) antibody (Peroxidase)

Species/Host Rabbit

Reactivity Mouse

Conjugation HRP

Tested Applications ELISA, IHC, WB

Immunogen Mouse IgG whole molecule

Preservatives None

Form/Appearance Lyophilized

Concentration 10.0 mg/mL

Storage Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

Note For research use only

Application notesThis product is designed for ELISA and western blotting. This product is also

suitable for multiplex analysis, including multicolor imaging, utilizing various

commercial platforms.

Isotype lgG

Clonality Polyclonal

Antibody Type Secondary Antibody

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: info@biorbyt.com Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558 **Biorbyt LLC.**

68 TW Alexander Drive, Durham, NC, 27713, United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558





Purity


This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Mouse IgG and Mouse Serum.

Dilution Range


ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:10,000

Expiration Date

12 months from date of receipt.



In vitro characterization of the LCL scaffold and the LCL-TbpA hybrid antigens. (A) Mouse sera were evaluated in a whole cell ELISA against both wild-type N. meningitidis M982 (Black) and M982 with TbpB knocked out (M982 Δ tbpB; Gray). Sera were assayed in triplicate from individual mice (4 to 5 mice per group) and averaged and displayed as mean +/- SEM. Significance was determined as a significant increase in titer compared with mice that received adjuvant alone by two-way ANOVA with Sidak's multiple comparison test for both wildtype and knockout data sets. **p \leq 0.01, ****p \leq 0.0001.



Whole-cell serum IgG ELISA titres of mice immunized with alum, TbpA or Hp7-L10 when captured by (A) heat-killed N. gonorrhoeae strain MS11 or (B) heat-killed N. meningitidis strain M982. Female mice were used in the lower genital tract gonococcal colonization, while male mice were challenged systemically with N. meningitidis. Significance was determined as a significant increase in titer compared with mice that received alum alone by ordinary one-way ANOVA with post-hoc analysis by Dunnett's multiple comparisons test. ****p 0.0001.