

Product Datasheet

Human IgG (H&L) Antibody Peroxidase Conjugated (orb347262)

Description	Human IgG (H&L) antibody (Peroxidase)
Species/Host	Mouse
Reactivity	Human
Conjugation	HRP
Tested Applications	ELISA, IHC, WB
Immunogen	Human IgG whole molecule
Preservatives	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Form/Appearance	Lyophilized
Concentration	1.0 mg/mL
Storage	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note	For research use only
Application notes	Anti-Human IgG Peroxidase conjugate has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.
Isotype	IgG
Clonality	Polyclonal


Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+441223859353) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+14159065211) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)

Antibody Type	Secondary Antibody
Purity	This product was prepared from monospecific polyclonal ascites by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-peroxidase, anti-Mouse Serum, Human IgG and Human Serum.
Dilution Range	ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:5,000
Expiration Date	12 months from date of receipt.



ELISA of purified Mouse Anti-Human IgG Antibody Peroxidase Conjugated tested against purified Human IgG HRP. Each well was coated in duplicate with 1.0 μg of Human IgG (p/n orb346219). The working dilution is 1:28000. The starting dilution of antibody was 5 $\mu\text{g}/\text{ml}$ and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate (p/n orb348651).

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)