

## Product Datasheet

### Mouse IgG (H&L) Secondary Antibody Fluorescein Conjugated Pre-Adsorbed (orb347377)

<b>Catalog Number</b>	orb347377
<b>Category</b>	Antibodies
<b>Description</b>	Mouse IgG (H&L) antibody (FITC)
<b>Clonality</b>	Polyclonal
<b>Species/Host</b>	Goat
<b>Isotype</b>	IgG
<b>Conjugation</b>	FITC
<b>Reactivity</b>	Mouse
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	1.0 mg/mL
<b>Buffer/Preservatives</b>	0.01% (w/v) Sodium Azide
<b>Purity</b>	Goat Anti-Mouse IgG was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.
<b>Immunogen</b>	Mouse IgG whole molecule
<b>Tested applications</b>	DOT, FC, FLISA, IF, WB

#### Biorbyt Ltd.

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom


Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

#### Biorbyt LLC


68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

<b>Dilution range</b>	FLISA: 1:10,000 - 1:50,000, FC: 1:500 - 1:2,500, IF: 1:1,000 - 1:5,000
<b>Application notes</b>	Anti-Mouse IgG Fluorescein Antibody has been tested by dot blot and western blot and is ideal for Fluorescent Western Blot, FLISA, Flow Cytometry, Immunohistochemistry and Immunofluorescence Microscopy as well as other antibody detection methods.
<b>Antibody Type</b>	Secondary Antibody
<b>Storage</b>	Store anti mouse secondary antibody at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note</b>	For research use only



Dot Blot showing the detection of Mouse IgG. A three-fold serial dilution of Mouse IgG starting at 200 ng was spotted onto 0.45 μm nitrocellulose. After blocking in 5% Blotto (p/n orb348624) 1 Hour at 20°C, Anti-Mouse IgG (H&L) (GOAT) Antibody Fluorescein Conjugated (p/n orb347377) secondary antibody was used at 1:1000 in Blocking Buffer for Fluorescent Western Blotting (p/n orb348637).



Western Blot of Goat anti-Mouse IgG Pre-Adsorbed Fluorescein Conjugated Secondary Antibody. Lane 1: Mouse IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Fluorescein goat secondary antibody at 1:1000 for 60 min at RT. Block: orb348637 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Mouse IgG. Other band(s): None.

#### Biorbyt Ltd.

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

#### Biorbyt LLC

68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558