

Product Datasheet

Mouse IgG (H&L) Secondary Antibody Peroxidase Conjugated Pre-Adsorbed (orb347433)

Catalog Number	orb347433
Description	MOUSE IgG (H&L) antibody (Peroxidase)
Species/Host	Goat
Reactivity	Mouse
Conjugation	HRP
Tested Applications	ELISA, IHC, WB
Immunogen	Mouse IgG whole molecule
Preservatives	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Form/Appearance	Lyophilized
Concentration	1.0 mg/mL
Storage	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note	For research use only
Application notes	Mouse secondary antibody conjugated to horseradish peroxidase is available in a variety of formats. Anti-Mouse IgG Peroxidase Antibody has been tested by ELISA and is suitable for ELISA, Immunohistochemistry western blotting as well as other HRP antibody based assays.
lsotype	lgG

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>


Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>



Biorbyt.com

Clonality	Polyclonal
Antibody Type	Secondary Antibody
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti- Peroxidase, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Human Serum Proteins.
Dilution Range	ELISA: 1:10,000-1:100,000, IHC: 1:500-1:5,000, WB: 1:5,000-1:40,000
Expiration Date	12 months from date of receipt.



ELISA results of purified Goat anti-Mouse IgG Antibody Peroxidase Conjugated (Min x Human Serum Proteins) tested against purified Mouse IgG. Each well was coated in duplicate with 1.0 μ g of Mouse IgG (p/n orb346219). The starting dilution of antibody was 5 μ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% fish gelatin and TME ELISA Peroxidase Substrate (p/n orb348651).

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>